

B²

[0188] Asparagine glycosylation sites have the following consensus pattern, N-{P}-[ST]-{P}, wherein N represents the glycosylation site. However, it is well known that that potential N-glycosylation sites are specific to the consensus sequence Asn-Xaa-Ser/Thr. However, the presence of the consensus tripeptide is not sufficient to conclude that an asparagine residue is glycosylated, due to the fact that the folding of the protein plays an important role in the regulation of N-glycosylation. It has been shown that the presence of proline between Asn and Ser/Thr will inhibit N-glycosylation; this has been confirmed by a recent statistical analysis of glycosylation sites, which also shows that about 50% of the sites that have a proline C-terminal to Ser/Thr are not glycosylated. Additional information relating to asparagine glycosylation may be found in reference to the following publications, which are hereby incorporated by reference herein: Marshall R.D., Annu. Rev. Biochem. 41:673-702(1972); Pless D.D., Lennarz W.J., Proc. Natl. Acad. Sci. U.S.A. 74:134-138(1977); Bause E., Biochem. J. 209:331-336(1983); Gavel Y., von Heijne G., Protein Eng. 3:433-442(1990); and Miletich J.P., Broze G.J. Jr., J. Biol. Chem. 265:11397-11404(1990).

On page 58, replace paragraph [0196] with the following paragraph:

B³

[0196] The putative consensus sequence for GPCRs comprises the conserved triplet and also spans the major part of the third transmembrane helix, and is as follows: [GSTALIVMFYWC]-[GSTANCPDE]-{EDPKRH}-x(2)-[LIVMNQGA]-x(2)-[LIVMFT]-[GSTANC]-[LIVMFYWSTAC]-[DENH]-R-[FYWCSH]-x(2)-[LIVM], where "X" represents any amino acid.